

SHARP TRANSITION BETWEEN EXTINCTION AND PROPAGATION OF REACTION

ANDREJ ZLATOŠ

ABSTRACT. We consider the reaction-diffusion equation

$$T_t = T_{xx} + f(T)$$

on $\mathbb{R}$ with $T_0(x) \equiv \chi_{[-L, L]}(x)$ and $f(0) = f(1) = 0$. In 1964 Kanel' proved that if f is an ignition non-linearity, then $T \rightarrow 0$ as $t \rightarrow \infty$ when $L < L_0$, and $T \rightarrow 1$ when $L > L_1$. We answer the open question of relation of L_0 and L_1 by showing that $L_0 = L_1$. We also determine the large time limit of T in the critical case $L = L_0$, thus providing the phase portrait for the above PDE with respect to a 1-parameter family of initial data. Analogous results for combustion and bistable non-linearities are proved as well.

1. INTRODUCTION

In the present paper we consider the reaction-diffusion equation

$$T_t = \Delta T + f(T) \tag{1.1}$$

in the cylinder $\mathbb{R} \times \Omega$ where Ω is a domain in $\mathbb{R}^{n-1}$, with Neumann boundary conditions on $\mathbb{R} \times \partial\Omega$. The non-linear reaction term f is assumed to be Lipschitz continuous with $f(0) = f(1) = 0$ and the initial datum T_0 is between 0 and 1.

We will treat the case when T_0 is independent of the transversal variable $y \in \Omega$, and so (1.1) becomes

$$T_t = T_{xx} + f(T) \tag{1.2}$$

with $x \in \mathbb{R}$. This equation has been extensively studied in mathematical, physical and other literature, starting with the pioneering works of Fisher [7] and Kolmogorov, Petrovskii, Piskunov [11]. In these papers (1.2) was used to describe the propagation of advantageous genes in a population. The main object of study in these and many subsequent works was the existence and stability of traveling fronts for (1.2) and (1.1). In the recent years most of the results have been extended to include an advection term $u \cdot \nabla T$ in (1.1), and we refer to the reviews [2, 16] for an extensive bibliography.

1991 *Mathematics Subject Classification*. Primary: 35K57; Secondary: 35K15 .

The above equations are used to model not only population genetics phenomena. When $f(\theta) > 0$ for $\theta \in (0, 1)$, then f is a *combustion non-linearity* and (1.1)/(1.2) model an exothermic chemical reaction in an infinite tube with a zero heat-loss boundary, in particular, flame propagation in a premixed combustible gas without advection (see Zel'dovich and Frank-Kamenetskii [17]). In this setting T is the normalized temperature taking values in $[0, 1]$. We note that (1.1) is usually obtained from a system involving both the temperature and the concentration of the reactants after the simplifying assumption of equal thermal and material diffusivities.

A special case of positive f , used often in chemical and biological literature, is the *KPP type* with $f''(\theta) \leq c < 0$ [11]. In combustion models the non-linearity is often considered to be of *Arrhenius type* with slow reaction rates at low temperatures, modeled by $f(\theta) = e^{-A/\theta}(1 - \theta)$. One often approximates this situation by considering an *ignition non-linearity* f satisfying $f(\theta) = 0$ for $\theta \in [0, \theta_0]$ and $f(\theta) > 0$ for $\theta \in (\theta_0, 1)$, with $\theta_0 \in (0, 1)$ the *ignition temperature*.

The third prominent case is the *bistable non-linearity* with $f(\theta) < 0$ for $\theta \in (0, \theta_0)$ and $f(\theta) > 0$ for $\theta \in (\theta_0, 1)$, where one usually assumes $\int_0^1 f(\theta) d\theta > 0$. This has been used to model signal propagation along bistable transmission lines, in particular, nerve pulse propagation [12]. In biological context it is also called *heterozygote inferior* (see Aronson and Weinberger [1]).

In this paper we will consider all the above types. Our interest here will not be in the question of traveling fronts, but in *extinction* of reaction — *quenching* of flames. We will therefore assume the initial datum $T_0(x)$ for (1.2) to be compactly supported, and will want to know when

$$\|T(t, \cdot)\|_\infty \rightarrow 0 \text{ as } t \rightarrow \infty. \quad (1.3)$$

For the sake of simplicity we will restrict ourselves to the case of T_0 being the characteristic function of an interval,

$$T_0(x) \equiv \chi_{[-L, L]}(x), \quad (1.4)$$

and study how long-time behavior of T depends on L . The methods in this paper allow one to treat some other increasing 1-parameter families of initial conditions, too.

Thus, we will study the competition of reaction and diffusion. The former helps increasing the temperature, whereas the latter (together with the compactness of the support of the initial datum) works towards the extinction of the flame. This question was originally addressed forty years ago by Kanel' [9] who considered the case of ignition

non-linearity and proved that if the initial datum is large enough, then reaction wins, whereas if it is small then diffusion manages to quench the flame. More precisely, when T solves (1.2)/(1.4) and f is of ignition type, Kanel' proved that there are two length scales L_0, L_1 such that

$$\begin{aligned} T(t, x) &\rightarrow 0 \text{ as } t \rightarrow \infty \text{ uniformly in } x \in \mathbb{R} \text{ if } L < L_0, \\ T(t, x) &\rightarrow 1 \text{ as } t \rightarrow \infty \text{ uniformly on compacts if } L > L_1. \end{aligned}$$

This has been extended to the case of bistable f by Aronson-Weinberger [1]. Both results actually hold when (1.4) is replaced by

$$T_0(x) \equiv \alpha \chi_{[-L, L]}(x). \quad (1.5)$$

for any $\alpha > \theta_0$, with L_0 and L_1 depending on α (in the ignition case this follows from [9], in the bistable case it was proved by Fife and McLeod [6]). A natural question arises: does L_0 equal L_1 ? If this is true and if one could determine the behavior of T as $t \rightarrow \infty$ when $L = L_0$, then one would be able to provide the complete “phase portrait” for the PDE (1.2) with respect to a 1-parameter family of initial conditions.

Since these early works, particularly in the recent years, several authors have studied quenching for (1.1). The above results have been extended to the case when (1.1) includes an advection term $u \cdot \nabla T$, with u a shear or periodic flow (see [13, 15]), even for certain combustion non-linearities [18]. Quenching of large initial data by large amplitude shear and cellular flows has been studied in [4, 5, 10, 18]. However, the question whether $L_0 = L_1$ remained open even in the simplest case of (1.2). The following two results provide the answer, including the treatment of the critical case $L = L_0$.

The first of them holds for ignition and combustion non-linearities.

Theorem 1. *Let $\theta_0 \in [0, 1)$ and $f : [0, 1] \rightarrow \mathbb{R}$ be Lipschitz with $f(\theta) = 0$ when $\theta \in [0, \theta_0]$, $f(\theta) > 0$ when $\theta \in (\theta_0, 1)$, and $f(1) = 0$. If $\theta_0 > 0$ then assume in addition that f is non-decreasing on $[\theta_0, \theta_0 + \delta]$ for some $\delta > 0$. Let $T : [0, \infty) \times \mathbb{R} \rightarrow [0, 1]$ solve*

$$\begin{aligned} T_t &= T_{xx} + f(T) \\ T_0(x) &\equiv \chi_{[-L, L]}(x). \end{aligned} \quad (1.6)$$

Then there is $L_0 \geq 0$ such that

- (i) *if $L < L_0$, then $T \rightarrow 0$ uniformly on $\mathbb{R}$ as $t \rightarrow \infty$;*
- (ii) *if $L = L_0$, then $T \rightarrow \theta_0$ uniformly on compacts as $t \rightarrow \infty$;*
- (iii) *if $L > L_0$, then $T \rightarrow 1$ uniformly on compacts as $t \rightarrow \infty$.*

Remark. The possibility of $L_0 = 0$ (so called *hair-trigger effect*) cannot be excluded when $\theta_0 = 0$. More precisely, using results from [18] one can show that $L_0 = 0$ when $f(\theta) \geq c\theta^p$ for some $p < 3$ and all

small θ , but $L_0 > 0$ when $f(\theta) \leq c\theta^p$ for some $p > 3$ and all small θ . We also note that if $\theta_0 = 0$, then the convergence in (ii) is as in (i) — uniform on $\mathbb{R}$.

Our second result holds for bistable non-linearities. We define $\theta_2 \in (\theta_0, 1)$ by $\int_0^{\theta_2} f(\theta)d\theta = 0$ and let U be the unique function solving $0 = U'' + f(U)$ with $U(0) = \theta_2$ and $U'(0) = 0$. Then U is an even function and we will show in the proof of the following theorem that it is positive on $\mathbb{R}$, decreasing to 0 on $(0, \infty)$, and bell-shaped.

Theorem 2. *Let $\theta_0 \in (0, 1)$ and $f : [0, 1] \rightarrow \mathbb{R}$ be Lipschitz with $f(0) = f(\theta_0) = f(1) = 0$, $f(\theta) < 0$ when $\theta \in (0, \theta_0)$, and $f(\theta) > 0$ when $\theta \in (\theta_0, 1)$. Assume also that $\int_0^1 f(\theta)d\theta > 0$. Let $T : [0, \infty) \times \mathbb{R} \rightarrow [0, 1]$ solve the problem (1.6). Then there is $L_0 > 0$ such that*

- (i) *if $L < L_0$, then $T \rightarrow 0$ uniformly on $\mathbb{R}$ as $t \rightarrow \infty$;*
- (ii) *if $L = L_0$, then $T \rightarrow U$ uniformly on $\mathbb{R}$ as $t \rightarrow \infty$;*
- (iii) *if $L > L_0$, then $T \rightarrow 1$ uniformly on compacts as $t \rightarrow \infty$.*

Remark. Both results can be extended to some other increasing families of initial conditions. In particular, to (1.5) with $\alpha > \theta_0$.

The crux of the proofs of both theorems will be to show that there is a single L for which T does not converge to 0 or 1 at $x = 0$ as $t \rightarrow \infty$. In Theorem 1 this will be achieved with the help of Lemma 4 by comparing solutions of (1.6) for two different initial conditions at differently rescaled times. In Theorem 2 it will follow from a detailed analysis of the large time behavior of T when the above limit is not 0 or 1, and an application of the comparison principle.

We note here that Theorem 1 is, in a sense, a limiting case of Theorem 2. If one takes $f \rightarrow 0$ on $(0, \theta_0)$ keeping f unchanged on $(\theta_0, 1)$, one has $\theta_2 \rightarrow \theta_0$ and $U \rightarrow \theta_0$ on compacts. That is, the bell shaped solution U from Theorem 2(ii) converges to the constant solution θ_0 from Theorem 1(ii).

The rest of the paper is devoted to the proofs of the two theorems. Section 2 contains preliminary Lemmas 3 and 4. Sections 3 and 4 prove Theorems 1 and 2, respectively.

The author thanks Henri Berestycki, Peter Constantin, François Hamel, Alexander Kiselev, Peter Poláčik, Jean-Michel Roquejoffre, and Lenya Ryzhik for encouragement and useful discussions. He also acknowledges the hospitality of the Mathematics Department of the University of Chicago where part of this work was done, as well as partial support from the NSF through the grant DMS-0314129.

2. PRELIMINARY LEMMAS

We start with

Lemma 3. *Let $f : [0, 1] \rightarrow \mathbb{R}$ be Lipschitz with $f(0) = f(1) = 0$. If $T : [0, \infty) \times \mathbb{R} \rightarrow [0, 1]$ solves (1.6), then the following hold.*

- (i) *If $|x| \leq |y|$ then $T(t, x) \geq T(t, y)$.*
- (ii) *There is $t_* > 0$ (possibly $t_* = \infty$) such that $T(t, 0)$ as a function of t is non-increasing on $[0, t_*)$ and non-decreasing on $[t_*, \infty)$.*
- (iii) *If $f \geq 0$ then there is $\theta_* \in [0, 1]$ such that $f(\theta_*) = 0$ and $T(t, x) \rightarrow \theta_*$ as $t \rightarrow \infty$, uniformly on compacts.*

Remarks. 1. For sufficiently smooth f this is essentially a result of Kanel' [9], although in (ii) he only proves $T(t, 0)$ to be eventually monotone.

2. In the case of (1.5) with $\alpha \in (\theta_0, 1)$, part (ii) has $0 < t_* \leq t_{**} \leq \infty$ such that $T(t, 0)$ is non-decreasing on $[0, t_*)$, non-increasing on $[t_*, t_{**})$ and non-decreasing on $[t_{**}, \infty)$.

Proof. We first assume that f is smooth and briefly recall main points of the proofs of (i) and (ii) from [9]. Let T^ε solve (1.6) but with initial condition $T(0, x) \equiv \chi^\varepsilon(x)$ where χ^ε are smooth, symmetric, decreasing in $|x|$, and converge to $\chi_{[-L, L]}$ in $L^1(\mathbb{R})$ as $\varepsilon \rightarrow 0$. Then $T_x^\varepsilon(0, x) \leq 0$ for $x > 0$, and by symmetry $T_x^\varepsilon(t, 0) = 0$. Since

$$(T_x^\varepsilon)_t = (T_x^\varepsilon)_{xx} + f'(T^\varepsilon)T_x^\varepsilon,$$

the maximum principle gives $T_x^\varepsilon(t, x) \leq 0$ for $x > 0$. Symmetry then yields $T_x^\varepsilon(t, x) \geq 0$ for $x < 0$. Since for any fixed $t > 0$ we have $T^\varepsilon(t, x) \rightarrow T(t, x)$ uniformly in x as $\varepsilon \rightarrow 0$, this proves (i).

Now let $D^h(t, x) \equiv T(t+h, x) - T(t, x)$. By the mean value theorem,

$$D_t^h = D_{xx}^h + f'(S)D^h$$

for some $S(t, x)$. Let Δ_h be set of (t, x) for which $D_h(t, x) \leq 0$. Then $\Delta_h \cap (\{0\} \times \mathbb{R}) = \{0\} \times [-L, L]$. By the maximum principle and symmetry, Δ_h is connected and its sections by lines parallel to the x -axis are segments symmetric about the t -axis. Therefore there is $0 < t_*^h \leq \infty$ such that $D^h(t, 0) < 0$ for $t \in [0, t_*^h)$ and $D^h(t, 0) \geq 0$ for $t \in [t_*^h, \infty)$. From $D^h(t, x) = D^{h/2}(t + \frac{h}{2}) + D^{h/2}(t)$ we obtain $t_*^{h/2} \in [t_*^h, t_*^h + \frac{h}{2}]$, and (ii) follows with $t_* \equiv \lim_{n \rightarrow \infty} t_*^{2^{-n}}$.

If f is only Lipschitz, take smooth f^ε such that $\|f^\varepsilon - f\|_\infty \leq \varepsilon$ and let T^ε solve (1.6) with f^ε in place of f . One can then show that $V^\varepsilon \equiv T^\varepsilon - T$ satisfies $|V^\varepsilon(t, x)| \leq \frac{\varepsilon}{c}(e^{ct} - 1)$ with c the Lipschitz constant for f (we spell this argument out in the proof of Theorem 1 below).

Therefore $T(t, x) = \lim_{\varepsilon \rightarrow 0} T^\varepsilon(t, x)$ for all t and x , and since (i) and (ii) hold for each T^ε , they also hold for T .

Finally, assume that $f \geq 0$. By (ii), $\theta_* \equiv \lim_{t \rightarrow \infty} T(t, 0)$ is well defined. Let Φ solve $\Phi_t = \Phi_{xx}$ on $\mathbb{R}^+$ with $\Phi(0, x) \equiv T(0, x)$ and boundary condition $\Phi(t, 0) \equiv T(t, 0)$. Then $\Phi(t, x) \rightarrow \theta_*$ as $t \rightarrow \infty$, uniformly on compacts. Since by the comparison principle (e.g., [14]) and (i), $\Phi(t, x) \leq T(t, x) \leq T(t, 0)$, the second claim in (iii) follows.

To prove the first claim, assume $f(\theta_*) > 0$ and choose $\varepsilon > 0$ such that for $\theta \leq \theta_* + 10\varepsilon$ we have $f(\theta) \geq \theta - \theta_* + 2\varepsilon$. Pick t_0 such that if Φ solves $\Phi_t = \Phi_{xx}$ on $\mathbb{R}$ with initial condition $\Phi(t_0, x) = T(t_0, x)$, then $\Phi(t, 0) \geq \theta_* - \varepsilon$ and $T(t, x) \leq \theta_* + \varepsilon$ for $t \in [t_0, t_0 + \ln 4]$ and $x \in \mathbb{R}$. This is possible thanks to the second claim in (iii). Define

$$S(t, x) \equiv \theta_* - 2\varepsilon + (\Phi(t, x) - \theta_* + 2\varepsilon)e^{t-t_0}.$$

Then $S(t, x) \leq \theta_* + 10\varepsilon$ for $t \in [t_0, t_0 + \ln 4]$ because $\Phi(t, x) \leq T(t, x) \leq \theta_* + \varepsilon$ for these t . A simple computation now shows that $S_t \leq S_{xx} + f(S)$ for $t \in [t_0, t_0 + \ln 4]$. Hence, S is a subsolution of (1.6) with $S(t_0, x) = T(t_0, x)$, and so $S \leq T$ for $t \in [t_0, t_0 + \ln 4]$. But $S(t_0 + \ln 4, 0) \geq \theta_* + 2\varepsilon > T(t_0 + \ln 4, 0)$, which is a contradiction. Therefore we have to have $f(\theta_*) = 0$. $\square$

Next, observe that we can use scaling to replace the variation in the initial condition in (1.6) by variation in the reaction strength. If T solves (1.6) with $T(0, x) \equiv \chi_{[-L, L]}(x)$, define $\tilde{T}(t, x) \equiv T(L^2 t, Lx)$, so that we have

$$\tilde{T}_t = \tilde{T}_{xx} + L^2 f(\tilde{T})$$

and $\tilde{T}(0, x) = \chi_{[-1, 1]}(x)$. Hence, Theorem 1 will be proved if we show that its conclusion holds for the L -dependent family of problems

$$T_t = T_{xx} + Lf(T) \tag{2.1}$$

$$T_0(x) \equiv \chi_{[-1, 1]}(x)$$

instead of (1.6) (note that Lemma 3 holds here, too). This important observation motivates the following key lemma.

Lemma 4. *Let Ω be a connected open domain in $\mathbb{R}^n$ with a smooth boundary (possibly $\Omega = \mathbb{R}^n$) and let $f, g : [0, \infty) \rightarrow \mathbb{R}$ be Lipschitz with $f(0) = g(0) = 0$ and $f \leq g$. Let $T, S : [0, \infty) \times \Omega \rightarrow [0, \infty)$ be continuous functions solving*

$$T_t = \Delta T + f(T) \tag{2.2}$$

$$S_t = \Delta S + g(S) \tag{2.3}$$

in Ω with Dirichlet boundary conditions on $\partial\Omega$. Assume $0 \leq T(0, x) \leq S(0, x)$ for all $x \in \Omega$ and $T(0, x_0) < S(0, x_0)$ for some x_0 . Assume also

that for any $\theta > 0$ the set $\Omega_{0,\theta} \equiv \{x \in \Omega \mid S(0, x) \geq \theta\}$ is compact. Finally, assume that there are $\theta_1 > 0$ and $\varepsilon_1 > 0$ such that for any $\theta \in [\theta_1, \|T\|_\infty]$ and $\varepsilon \in [0, \varepsilon_1]$ we have

$$g(\theta + \varepsilon[\theta - \theta_1]) \geq (1 + \varepsilon)f(\theta). \quad (2.4)$$

Then

$$\liminf_{t \rightarrow \infty} \inf_{T(t,x) > \theta_1} \frac{S(t, x) - \theta_1}{T(t, x) - \theta_1} > 1 \quad (2.5)$$

with the convention that infimum over an empty set is ∞ .

Remark. The result holds without change when we add a first order term $u(x) \cdot \nabla$ with $u \in C^1$ to (2.2) and (2.3).

Proof. First notice that the assumptions imply that

$$\Omega_{t,\theta} \equiv \{x \in \Omega \mid S(t, x) \geq \theta\}$$

are compact. Indeed, by the maximum principle, $\Omega_{t,\theta} \subseteq \tilde{\Omega}_{t,\delta\theta}$ where $\delta \equiv e^{-tc}$ with c the Lipschitz constant for g , and $\tilde{\Omega}_{t,\theta}$ is defined as $\Omega_{t,\theta}$ but with Φ , the solution of

$$\Phi_t = \Delta \Phi, \quad \Phi(0, x) = S(0, x),$$

in place of S . Compactness of $\tilde{\Omega}_{t,\theta}$ follows from that of $\tilde{\Omega}_{0,\theta}$ and the Feynman-Kac formula.

The assumptions and the strong maximum principle also imply $T(t, x) < S(t, x)$ for $t > 0$ and $x \in \Omega$. Let us define

$$\begin{aligned} \Omega(t) &\equiv \{x \in \Omega \mid T(t, x) > \theta_1\}, \\ \Omega'(t) &\equiv \{x \in \Omega \mid T(t, x) = \theta_1\}, \end{aligned}$$

and let

$$\omega(t) \equiv \min \left\{ 1 + \varepsilon_1, \inf_{x \in \Omega(t)} \frac{S(t, x) - \theta_1}{T(t, x) - \theta_1} \right\}.$$

Since $\overline{\Omega(t)}$ is compact and $T < S$ continuous, $\omega(t) > 1$ for $t > 0$. Hence the result will follow if we show that ω is a non-increasing function. Since ω is continuous (because T and S are), it is sufficient to show that for any $t_0 > 0$ there is $\tau_0 > 0$ such that for all $t \in [t_0, t_0 + \tau_0]$ we have $\omega(t) \geq \omega(t_0)$.

Hence, fix $t_0 > 0$. Notice that uniform boundedness of the sets Ω_{t,θ_1} for $t \in [0, t_0 + 1]$ (by the maximum principle and Feynman-Kac formula) and continuity of T, S imply that T, S are uniformly continuous on $[0, t_0 + 1] \times \Omega$, and the set

$$\Sigma \equiv \{(t, x) \mid t \in [t_0, t_0 + 1] \text{ and } x \in \Omega(t) \cup \Omega'(t)\}$$

is compact. Thanks to the uniform continuity of T we only need to consider the case $\Omega(t_0) \cup \Omega'(t_0) \neq \emptyset$, and hence $\Sigma \neq \emptyset$. Since S is continuous and $S > T$,

$$\sigma \equiv \inf_{(t,x) \in \Sigma} \{S(t,x) - \theta_1\} > 0.$$

We let $\delta \equiv \sigma/4(1 + \varepsilon_1)$ and define

$$\Delta \equiv \{x \mid |T(t_0, x) - \theta_1| \leq \delta \text{ and } S(t_0, x) - \theta_1 \geq \sigma - \delta\}$$

By the uniform continuity of T, S , there is $\tau_0 \in (0, 1)$ such that for $t \in [t_0, t_0 + \tau_0]$ and $x \in \Omega$ we have

$$|T(t, x) - T(t_0, x)| \leq \frac{\delta}{2} \quad \text{and} \quad |S(t, x) - S(t_0, x)| \leq \frac{\delta}{2}. \quad (2.6)$$

So if $t \in [t_0, t_0 + \tau_0]$, then

$$\Omega(t) \subseteq \Omega(t_0) \cup \Delta \quad (2.7)$$

(note that $S(t_0, x) - \theta_1 \geq \sigma - \frac{\delta}{2}$ for $x \in \Omega(t)$ because then $(t, x) \in \Sigma$). Now if $t \in [t_0, t_0 + \tau_0]$ and $x \in \Delta$, then by (2.6),

$$S(t, x) - \theta_1 > \frac{\sigma}{2} > (1 + \varepsilon_1)|T(t, x) - \theta_1| \geq \omega(t_0)(T(t, x) - \theta_1). \quad (2.8)$$

Next let

$$A \equiv \{x \in \Omega \mid T(t_0, x) > \theta_1 + \delta\} = \Omega(t_0) \setminus \Delta \quad (2.9)$$

and

$$B \equiv \{x \in \Omega \mid T(t_0, x) \geq \theta_1 + \frac{\delta}{2}\} \subseteq \Omega(t_0).$$

Uniform continuity of T shows that, $\text{dist}(A, B^c) > 0$, and so there is an open set Γ with a smooth boundary such that $A \subseteq \Gamma \subseteq B$. Let $\tilde{T} \equiv T - \theta_1$, $\tilde{U} \equiv \omega(t_0)\tilde{T}$, $\tilde{S} \equiv S - \theta_1$, $\tilde{f}(\theta) \equiv f(\theta + \theta_1)$, and $\tilde{g}(\theta) \equiv g(\theta + \theta_1)$. Then for $x \in \Gamma$ we have

$$\tilde{S}(t_0, x) \geq \omega(t_0)\tilde{T}(t_0, x) = \tilde{U}(t_0, x)$$

by the definition of $\omega(t_0)$, for $t \in [t_0, t_0 + \tau_0]$ and $x \in \partial\Gamma$ we have

$$\tilde{S}(t, x) > \sigma - 2\delta > \frac{\sigma}{2} \geq \omega(t_0)\tilde{T}(t, x) = \tilde{U}(t, x)$$

since $\partial\Gamma \subseteq B \setminus A \subseteq \Delta$, and for $t \in [t_0, t_0 + \tau_0]$ and $x \in \Gamma$ we have

$$\begin{aligned} \tilde{U}_t &= \Delta \tilde{U} + \omega(t_0)\tilde{f}\left(\frac{1}{\omega(t_0)}\tilde{U}\right), \\ \tilde{S}_t &= \Delta \tilde{S} + \tilde{g}(\tilde{S}) \end{aligned}$$

by (2.2) and (2.3). For these (t, x) we have $T(t, x) \geq \theta_1$ because of (2.6) and $\Gamma \subseteq B$, and so by (2.4) and $\omega(t_0) - 1 \in (0, \varepsilon_1]$,

$$\omega(t_0)\tilde{f}\left(\frac{1}{\omega(t_0)}\tilde{U}\right) = \omega(t_0)f(T) \leq g(\omega(t_0)[T - \theta_1] + \theta_1) = \tilde{g}(\tilde{U}).$$

The comparison principle now shows that $\tilde{S} \geq \tilde{U}$ on $[t_0, t_0 + \tau_0] \times \Gamma$. Hence

$$S(t, x) - \theta_1 \geq \omega(t_0)(T(t, x) - \theta_1)$$

for $t \in [t_0, t_0 + \tau_0]$ and $x \in A$, which together with (2.8), (2.9), and (2.7) gives $\omega(t) \geq \omega(t_0)$ for $t \in [t_0, t_0 + \tau_0]$. The proof is finished. $\square$

3. PROOF OF THEOREM 1

We can now complete the proof of Theorem 1. We will do this for the formulation in (2.1).

First assume $\theta_0 > 0$. We know from Lemma 3(iii) that for every L we have $T \rightarrow \theta_*^L$ uniformly on compacts, with θ_*^L such that $f(\theta_*^L) = 0$. Obviously $\theta_*^L \notin (0, \theta_0)$ because in that case we would have $T(t, x) \leq \theta_0$ for all $t \geq t_0$ and consequently $T \rightarrow 0$ (since $\|T(t_0, \cdot)\|_1 < \infty$ and $T_t = T_{xx}$ for $t \geq t_0$). So we are only left with $\theta_*^L \in \{0, \theta_0, 1\}$.

Let A , B , and C be the sets of $L \geq 0$ such that θ_*^L equals 0, θ_0 , and 1, respectively. Notice that since $T(t, 0) \geq T(t, x)$, the convergence of T to 0 for $L \in A$ is actually uniform on $\mathbb{R}$. We have $A \cup B \cup C = [0, \infty)$ and the comparison principle implies that the three sets are intervals with A lying to the left of B and B to the left of C .

Moreover, A and C are non-empty by Kanel' [9] and open. The latter follows from the fact that if T^L is the solution of (2.1), then for $L_1 < L_2$ and $V \equiv T^{L_2} - T^{L_1}$ we have $V \geq 0$ by comparison, and

$$\begin{aligned} V_t &= \Delta V + (L_2 - L_1)f(T^{L_2}) + L_1[f(T^{L_2}) - f(T^{L_1})] \\ &\leq \Delta V + c(L_2 - L_1) + cL_1V \end{aligned}$$

with $c \geq \|f\|_\infty$ the Lipschitz constant for f . Since the function $\tilde{V}(t, x) \equiv \frac{L_2 - L_1}{L_1}(e^{cL_1 t} - 1)$ satisfies

$$\tilde{V} = \Delta \tilde{V} + c(L_2 - L_1) + cL_1 \tilde{V}$$

with $\tilde{V}(0) = 0 = V(0)$, the comparison principle gives $V \leq \tilde{V}$, that is,

$$T^{L_2}(t, x) - T^{L_1}(t, x) \in [0, \frac{L_2 - L_1}{L_1}(e^{cL_1 t} - 1)].$$

Therefore if $L_1 \in A$, then $T^{L_1}(t_0, 0) \leq \frac{1}{2}\theta_0$ for some $t_0 > 0$, and hence $T^{L_2}(t_0, 0) < \theta_0$ (and so $L_2 \in A$) for $L_2 < L_1 + \frac{1}{2}\theta_0 L_1(e^{cL_1 t_0} - 1)^{-1}$. On the other hand, Kanel's result [9] also holds for (1.5), and it says that for any $\alpha > \theta_0$ and $L > 0$ there is $M = M(\alpha, L) < \infty$ such that if T solves (2.1) and $T(t_0, x) \geq \alpha \chi_{[-M, M]}(x)$, then $T \rightarrow 1$ uniformly on compacts. Let $\theta_0 < \alpha < \beta < 1$ and if $L_1 \in C$, let $M = M(\alpha, \frac{1}{2}L_1)$. For some t_0 we have $T^{L_1}(t_0, x) \geq \beta \chi_{[-M, M]}(x)$ and so for any $L_2 > L_1 - (\beta - \alpha)L_1(e^{cL_1 t_0} - 1)^{-1}$ we have $T^{L_2}(t_0, x) \geq \alpha \chi_{[-M, M]}(x)$. If in

addition $L_2 > \frac{1}{2}L_1$, we have $L_2 \in C$. So A, C are non-empty and open, and hence B is non-empty and closed.

The proof will be finished if we show that B contains a single element. Hence assume $L_1 < L_2$ are both in B . Let $\theta_1 \equiv \frac{1}{2}\theta_0 \in (0, \theta_0)$ and

$$\varepsilon_1 \equiv \min\{L_2 L_1^{-1} - 1, \delta(\delta + \theta_0)^{-1}\} > 0$$

(with δ from the statement of Theorem 1). Choose $t_0 > 0$ such that $T^{L_1}(t, x) \leq \theta_0 + \frac{\delta}{2}$ when $t \geq t_0$. The comparison principle, $f \not\equiv 0$, and the strong maximum principle yield $T^{L_1} < T^{L_2}$ for $t > 0$ and both T^{L_1} and T^{L_2} are obviously continuous for $t > 0$. Lipschitzness of f and compact support of $T^{L_2}(0, \cdot)$ show that for any $\theta > 0$, the set of x for which $T^{L_2}(t_0, x) \geq \theta$ is compact. Finally, whenever $\theta \in [\theta_1, \theta_0 + \frac{\delta}{2}]$ and $\varepsilon \in [0, \varepsilon_1]$, we have $\theta + \varepsilon[\theta - \theta_1] \leq \theta_0 + \delta$. Thus by the assumptions on f (and the definition of ε_1),

$$L_2 f(\theta + \varepsilon[\theta - \theta_1]) \geq L_2 f(\theta) \geq (1 + \varepsilon)L_1 f(\theta).$$

Therefore Lemma 4 applies to T^{L_1} and T^{L_2} (with starting time t_0) and shows that for some $r > 1$ and all large enough t we have

$$T^{L_2}(t, x) - \theta_1 \geq r[T^{L_1}(t, x) - \theta_1]$$

whenever $T^{L_1}(t, x) > \theta_1$. But since $\theta_1 < \theta_0$, this contradicts the assumption that both $T^{L_1}(t, 0)$ and $T^{L_2}(t, 0)$ converge to θ_0 as $t \rightarrow \infty$. Hence, $B = \{L_0\}$ and we are done.

Now, consider $\theta_0 = 0$. We have $\theta_*^L \in \{0, 1\}$, the sets A, C satisfy $A \cup C = [0, \infty)$, and by the comparison principle, A lies to the left of C . Moreover, $0 \in A$ and C is non-empty and open by the same argument as above. Hence A is closed and its maximum is L_0 (possibly $L_0 = 0$). Lemma 3(iii) yields (iii) of this theorem and $T(t, 0) \geq T(t, x)$ gives (i) and (ii), including the fact that the convergence in (ii) is uniform on $\mathbb{R}$. The proof is finished.

4. PROOF OF THEOREM 2

The situation is somewhat more complicated here. Firstly, we do not have Lemma 3(iii) at our disposal, and so the limit of T as $t \rightarrow \infty$ need not always be a constant function. And secondly, we cannot use Lemma 4 and the scaling argument preceding it in the way we did in the last section because it is not anymore true that $L_2 f \geq L_1 f$ when $L_2 > L_1$. We note that one can still use the lemma without scaling, but then the argument applies only to a restricted class of bistable f . Fortunately, it turns out that the first of these difficulties actually cancels the problems created by the second, as we shall see below.

Let us therefore go back to T solving (1.6) rather than (2.1). We know from Lemma 3(ii) that $\theta_*^L \equiv \lim_{t \rightarrow \infty} T(t, 0)$ is well defined, and from the comparison principle that it is non-decreasing in L .

First assume $\theta_*^L < \theta_2$, with θ_2 defined in the introduction. Choose $\varepsilon > 0$ and a Lipschitz function $\tilde{f} : [0, 1] \rightarrow \mathbb{R}$ so that $\tilde{f} = 0$ on $[0, \varepsilon]$, $\tilde{f}'(\varepsilon) < 0$, $\tilde{f} \geq f$ on $(\varepsilon, \frac{1}{2}(\theta_*^L + \theta_2)]$ and $\tilde{f}$ has a single zero there, $\tilde{f} > 0$ on $(\frac{1}{2}(\theta_*^L + \theta_2), 1)$, $\tilde{f}(1) = 0 > \tilde{f}'(1)$, and

$$\int_0^1 \tilde{f}(\theta) d\theta < 0. \quad (4.1)$$

Let t_0 be such that for $t \geq t_0$ and all $x \in \mathbb{R}$ we have $T(t, x) \leq \frac{1}{2}(\theta_*^L + \theta_2)$. This is possible by Lemma 3(i). Since $\tilde{f} \geq f$ on $[0, \frac{1}{2}(\theta_*^L + \theta_2)]$, starting from time t_0 one has $T_t \leq T_{xx} + \tilde{f}(T)$, that is, T is a subsolution of the equation

$$\Phi_t = \Phi_{xx} + \tilde{f}(\Phi). \quad (4.2)$$

Let $\phi : \mathbb{R} \rightarrow [0, 1]$ with $\phi(x) \rightarrow \varepsilon$ as $x \rightarrow \infty$ and $\phi(x) \rightarrow 1$ as $x \rightarrow -\infty$ be the unique, up to translation, traveling front profile (with speed v) for (4.2) [8]. That is, $\phi(x - vt)$ solves (4.2). It follows from (4.1) that in this case $v < 0$.

From compactness of the support of $T(0, x)$ and Lipschitzness of f , $T(t_0, x) \rightarrow 0$ as $|x| \rightarrow \infty$. This and $\|T(t_0, \cdot)\|_\infty < 1$ mean that there is x_0 such that $T(t_0, x) \leq \phi(x - x_0 - vt_0)$, and since $\phi(x - x_0 - vt)$ is a solution and $T(t, x)$ a subsolution of (4.2),

$$T(t, x) \leq \phi(x - x_0 - vt)$$

for all $t \geq t_0$. But then $T(t, 0) \leq \phi(-x_0 - vt) \rightarrow \varepsilon$ as $t \rightarrow \infty$ because $v < 0$. This holds for any small enough $\varepsilon > 0$ and thus $\theta_*^L = 0$.

Next assume $\theta_*^L > \theta_2$. Let S be the solution of (1.6) on $\mathbb{R}^+$ with $S(0, x) = 0$ and $S(t, 0) = s(t)$ a smooth strictly increasing function with all derivatives bounded such that $s(0) = 0$, $s(t) \leq T(t, 0)$, and $\lim_{t \rightarrow \infty} s(t) = \theta_*^L$. Then for any $h > 0$ we have $S(h, x) > S(0, x)$ and so by comparison $S(t + h, x) > S(t, x)$. Hence $\tilde{S}(x) \equiv \lim_{t \rightarrow \infty} S(t, x) > 0$ is well defined and $\tilde{S}(0) = \theta_*^L$. Since by comparison again $S(t, x) \leq T(t, x) \leq T(t, 0)$, we also have $\tilde{S}(x) \leq \theta_*^L$.

Standard parabolic regularity shows that $S(t, x)$ converges to $\tilde{S}(x)$ along with its first two derivatives uniformly on compacts, and so $\tilde{S}$ solves the stationary problem

$$0 = \tilde{S}'' + f(\tilde{S}) \quad (4.3)$$

on $\mathbb{R}^+$ (this can be found also in [1]). But then for any $y > 0$

$$\int_{\tilde{S}(y)}^{\theta_*^L} f(\theta) d\theta = \int_y^0 f(\tilde{S}(x)) \tilde{S}'(x) dx = \int_0^y \tilde{S}''(x) \tilde{S}'(x) dx = \frac{1}{2} [(\tilde{S}'(y))^2 - (\tilde{S}'(0))^2].$$

Assume there is $y > 0$ such that $\tilde{S}(y) < \theta_*^L$, and then pick one such that also $\tilde{S}'(y) < 0$. Since $\int_{\omega}^{\theta_*^L} f(\theta) d\theta$ is bounded below by a positive constant for all $\omega \in [0, \tilde{S}(y)]$, so is $-\tilde{S}'(z)$ for all $z \geq y$ (because $\tilde{S}(z) \geq 0$). But that means $\tilde{S}$ is eventually negative, a contradiction. Hence, we must have $\tilde{S} \equiv \theta_*^L$, which is only possible if $\theta_*^L = 1$. Moreover, since S converges to $\tilde{S} \equiv 1$ uniformly on compacts as $t \rightarrow \infty$ (and $S \leq T \leq 1$), so does T .

The above shows that $\theta_*^L \in \{0, \theta_2, 1\}$. As in the proof of Theorem 1, and using the equivalent of Kanel's result for (1.5) and bistable f [6], one can show that the intervals A, C of L for which $\theta_*^L = 0, 1$, respectively, are non-empty and open. If B is the interval of L for which $\theta_*^L = \theta_2$, then B lies between A and C and again $A \cup B \cup C = [0, \infty)$.

Next we need to prove that B only contains one element. We will show below that if $L \in B$, then $T(t, x) \rightarrow U(x)$ uniformly on $\mathbb{R}$ as $t \rightarrow \infty$. Here U solves (4.3) with $U(0) = \theta_2$ and $U'(0) = 0$. Assume now that $L_1 < L_2$ are both in B , with T^{L_1} and T^{L_2} the corresponding solutions of (1.6). We then have $T^{L_1}(t, 0) \rightarrow \theta_2$, and since the equation is translation invariant, we also have $\tilde{T}(t, \varepsilon) \rightarrow \theta_2$ when $\tilde{T}$ solves (1.6) with initial condition $\tilde{T}_0(x) \equiv \chi_{[-L_1+\varepsilon, L_1+\varepsilon]}(x)$. But if $|\varepsilon| < L_2 - L_1$, then $T_0^{L_2}(x) \geq \tilde{T}_0(x)$, and so by the comparison principle,

$$U(\varepsilon) = \lim_{t \rightarrow \infty} T^{L_2}(t, \varepsilon) \geq \theta_2.$$

Since $U''(0) = -f(U(0)) = -f(\theta_2) < 0$, U has a strict local maximum at zero and therefore $U(\varepsilon) < U(0) = \theta_2$ for all small enough $|\varepsilon| > 0$. This is a contradiction and hence $B = \{L_0\}$.

To complete the proof, we need to show that $T(t, x) \rightarrow U(x)$ uniformly as $t \rightarrow \infty$ when $L \in B$ (and hence $\theta_*^L = \theta_2$). Notice that the argument following (4.3) applies to U and we find for any $x > 0$ such that $U(x) \geq 0$,

$$\int_{U(x)}^{\theta_2} f(\theta) d\theta = \frac{1}{2} (U'(x))^2. \quad (4.4)$$

The definition of θ_2 then shows that $U(x) \leq \theta_2$, and $U'(x) \neq 0$ when $U(x) \in (0, \theta_2)$. Since $U(x)$ cannot be constant θ_2 on any interval and U' is continuous, we must have $U'(x) < 0$ for all $x > 0$ such that $U(x) > 0$. There is no x with $U(x) = 0$ because then (4.4) would give $U'(x) = 0$, contradicting uniqueness of solutions to initial value

problems associated to (4.3). Hence $U(x) \in (0, \theta_2)$ and $U'(x) < 0$ for $x > 0$, with $U'(x)$ bounded away from zero when $U(x)$ is away from zero (by (4.4) and the definition of θ_2). This and symmetry show that U is indeed a symmetric bell-shaped solution (with U' decreasing on $[0, U^{-1}(\theta_0)]$ and increasing on $[U^{-1}(\theta_0), \infty)$ by (4.4)) of the stationary problem (4.3) such that $U(x) \rightarrow 0$ as $|x| \rightarrow \infty$.

If we now apply the argument involving S and $\tilde{S}$ from the case $\theta_*^L > \theta_2$, we find as above that $\tilde{S}(x) \leq \theta_2 = \tilde{S}(0)$ and that $\tilde{S} > 0$ is possible only if $\tilde{S}'(0) = 0$. But then $\tilde{S}(0) = U(0)$ and $\tilde{S}'(0) = U'(0)$, thus $\tilde{S} = U$. Moreover, uniform on compacts convergence of S to U and $0 \leq S(t, x) \leq U(x) \rightarrow 0$ as $|x| \rightarrow \infty$ yield uniform on $\mathbb{R}$ convergence of S to U . Since $T(t, x) \geq S(t, x)$, we have $\liminf_{t \rightarrow \infty} T(t, x) \geq U(x)$ uniformly on $\mathbb{R}$. Hence we are left with proving $\limsup_{t \rightarrow \infty} T(t, x) \leq U(x)$ uniformly in $x > 0$ (which suffices due to symmetry).

Let $0 < x_0 < \infty$ be such that if $S(0, x) \geq \theta_2 \chi_{[-x_0, x_0]}(x)$ and S satisfies (1.6), then $S \rightarrow 1$ uniformly on compacts. Such x_0 exists by [6] because $\theta_2 > \theta_0$. Then obviously for every $t \geq 0$ we have $T(t, x_0) \leq \theta_2$, because otherwise Lemma 3(i) would imply $T \rightarrow 1$ uniformly on compacts. Since both $T(t, x)$ and $V(x) = U(x - x_0)$ satisfy (1.6) on (x_0, ∞) , $V(x_0) = \theta_2 \geq T(t, x_0)$, and $V(x) > 0 = T(0, x)$ for $x > x_0$, the comparison principle implies $T(t, x) \leq V(x)$. Let us therefore define

$$x_1 \equiv \min\{\tilde{x} \mid \limsup_{t \rightarrow \infty} T(t, x) \leq U(x - \tilde{x}) \text{ uniformly in } x > \tilde{x}\} \leq x_0.$$

The minimum is achieved because U is uniformly continuous and by Lemma 3(i), $\limsup_{t \rightarrow \infty} T(t, x) \leq \theta_2 = U(0)$. We note that at this point one can derive $T \rightarrow U$ from $x_1 < \infty$ and the results of [3] when $f \in C^1$ and $f'(0) < 0$. Our non-linearity is more general and so [3] is not applicable here.

If $x_1 = 0$, then we are done, so assume $x_1 > 0$. First notice that $\limsup_{t \rightarrow \infty} T(t, \frac{1}{2}x_1) \leq \theta_2 - \delta_1$ for some $\delta_1 > 0$. Indeed — in view of $\lim_{t \rightarrow \infty} T(t, 0) = \theta_2$, Lemma 3(i), and the comparison principle, it is sufficient to show that there are $\delta_1, \delta_2 > 0$ such that if $S(0, x) \geq (\theta_2 - \delta_1)\chi_{[-x_1/2, x_1/2]}(x)$ and S satisfies (1.6), then $S(t, 0) \geq \theta_2 + \delta_2$ for some $t > 0$. This in turn is true because it holds for $\delta_1 = 0$ and some $t, \delta_2 > 0$, since then $S(0, 0) = \theta_2$ and $S_t(0, 0) = f(\theta_2) > 0$, and because $S(t, 0)$ is continuous in δ_1 .

Now choose $x_2 \in (\frac{1}{2}x_1, x_1)$ such that $U(x_1 - x_2) \geq \theta_2 - \delta_1$. The above and Lemma 3(i) show that $\limsup_{t \rightarrow \infty} T(t, x) \leq \theta_2 - \delta_1$ uniformly in $x \geq \frac{1}{2}x_1$, and so

$$\limsup_{t \rightarrow \infty} T(t, x) \leq U(x - x_2) \tag{4.5}$$

uniformly in $x \in [x_2, x_1]$. We will show that (4.5) also holds uniformly in $x > x_1$, which will yield $x_1 \leq x_2$ by the definition of x_1 . This will be a contradiction and hence necessarily $x_1 = 0$.

Let $s(t)$ be smooth, decreasing with all derivatives bounded, such that $s(0) = \theta_2$ and $\lim_{t \rightarrow \infty} s(t) = U(x_1 - x_2)$. Let $S(t, x)$ satisfy (1.6) for $x > x_1$ with $S(0, x) = U(x - x_1)$ and $S(t, x_1) = s(t)$. As above, one proves that this time S is non-increasing in t ,

$$S(t, x) \in [U(x - x_2), U(x - x_1)], \quad (4.6)$$

and $\tilde{S}(x) \equiv \lim_{t \rightarrow \infty} S(t, x)$ satisfies (4.3) and $\tilde{S}(x) \rightarrow 0$ as $x \rightarrow \infty$ (by (4.6)). Moreover, $S \rightarrow \tilde{S}$ uniformly on compacts, which together with (4.6) and $U(x) \rightarrow 0$ as $x \rightarrow \infty$ shows that $S \rightarrow \tilde{S}$ uniformly on $\mathbb{R}$. Since $\tilde{S}(x_1) = U(x_1 - x_2)$ and $\tilde{S}(\infty) = U(\infty) = 0$, a formula similar to (4.4), with the integral from 0 to $U(x_1 - x_2)$, gives $\tilde{S}'(x_1) = U'(x_1 - x_2)$. It follows that $\tilde{S}(x) = U(x - x_2)$.

Now pick any $\varepsilon > 0$ and choose t_0 such that

$$S(t, x) - U(x - x_2) < \frac{\varepsilon}{2} \quad (4.7)$$

for $t \geq t_0$ and $x \geq x_1$. Then pick t_1 so that

$$T(t, x) - U(x - x_1) < \varepsilon_0 \equiv \frac{\varepsilon}{2} e^{-ct_0}$$

for $t \geq t_1$ and $x \geq x_1$ (with c the Lipschitz constant for f). This is possible by the definition of x_1 .

For any $t_2 > t_1$ and $x > x_1$ we have

$$T(t_2, x) - S(0, x) = T(t_2, x) - U(x - x_1) < \varepsilon_0,$$

and for $t > t_2$ we have

$$T(t, x_1) - S(t - t_2, x_1) = T(t, x_1) - s(t) \leq T(t, x_1) - U(x_1 - x_2) < \varepsilon_0$$

by (4.5) if t_2 is large enough. Hence if we let $R(t, x) \equiv S(t, x) + e^{ct}\varepsilon_0$, then $T(t_2, x) < R(0, x)$ for $x > x_1$, $T(t, x_1) < R(t - t_2, x_1)$ for $t > t_2$, and

$$R_t = S_t + ce^{ct}\varepsilon_0 = S_{xx} + f(S) + ce^{ct}\varepsilon_0 \geq R_{xx} + f(R).$$

So R is a supersolution of (1.6), and by the comparison principle $T(t, x) \leq R(t - t_2, x)$ for $t > t_2$ and $x > x_1$. In particular,

$$T(t_2 + t_0, x) \leq R(t_0, x) = S(t_0, x) + \frac{\varepsilon}{2} < U(x - x_2) + \varepsilon$$

for $x > x_1$ by (4.7). Since this holds for any large enough t_2 , we have $T(t, x) < U(x - x_2) + \varepsilon$ for all large t and $x > x_1$. As $\varepsilon > 0$ was arbitrary, this gives (4.5) uniformly in $x > x_1$. Hence $x_1 \leq x_2 < x_1$, a contradiction. Therefore we must have $x_1 = 0$ and the proof is finished.

REFERENCES

- [1] D.G. Aronson and H.F. Weinberger, *Nonlinear diffusion in population genetics, combustion, and nerve pulse propagation*, Partial differential equations and related topics, Lecture Notes in Math. **446**, Springer, Berlin, 1975.
- [2] H. Berestycki, *The influence of advection on the propagation of fronts in reaction-diffusion equations*, Nonlinear PDEs in Condensed Matter and Reactive Flows, NATO Science Series C, 569, H. Berestycki and Y. Pomeau eds, Kluwer, Dordrecht, 2003.
- [3] J. Busca, M.A. Jendoubi, and P. Poláčik, *Convergence to equilibrium for semi-linear parabolic problems in $\mathbb{R}^N$* , Comm. Partial Differential Equations **27** (2002), 1793–1814.
- [4] P. Constantin, A. Kiselev, and L. Ryzhik, *Quenching of flames by fluid advection*, Comm. Pure Appl. Math. **54** (2001), 1320–1342.
- [5] A. Fannjiang, A. Kiselev, and L. Ryzhik, *Quenching of reaction by cellular flow*, preprint.
- [6] P.C. Fife and J.B. McLeod, *The approach of solutions of nonlinear diffusion equations to travelling front solutions*, Arch. Ration. Mech. Anal. **65** (1977), 335–361.
- [7] R.A. Fisher, *The advance of advantageous genes*, Ann. of Eugenics **7** (1937), 355–369.
- [8] Ja.I. Kanel', *Stabilization of solutions of the Cauchy problem for equations encountered in combustion theory*, Mat. Sb. (N.S.) **59** (101) 1962 suppl., 245–288.
- [9] Ja.I. Kanel', *Stabilization of the solutions of the equations of combustion theory with finite initial functions*, Mat. Sb. (N.S.) **65** (107) 1964 suppl., 398–413.
- [10] A. Kiselev and A. Zlatoš, *Quenching of combustion by shear flows*, preprint.
- [11] A.N. Kolmogorov, I.G. Petrovskii, and N.S. Piskunov, *Étude de l'équation de la chaleur de matière et son application à un problème biologique*, Bull. Moskov. Gos. Univ. Mat. Mekh. **1** (1937), 1–25.
- [12] J. Nagumo, S. Arimoto, and S. Yoshizawa, *An active pulse transmission line simulating nerve axon*, Proc. Inst. Radio Eng. **50** (1962), 2061–2070.
- [13] J.-M. Roquejoffre, *Eventual monotonicity and convergence to travelling fronts for the solutions of parabolic equations in cylinders*, Ann. Inst. H. Poincaré Anal. Non Linéaire **14** (1997), 499–552.
- [14] J. Smoller, *Shock Waves and Reaction-Diffusion Equations*, Springer-Verlag, New York, 1994.
- [15] J. Xin, *Existence and nonexistence of travelling waves and reaction-diffusion front propagation in periodic media*, J. Stat. Phys. **73** (1993), 893–926.
- [16] J. Xin, *Front propagation in heterogeneous media*, SIAM Rev. **42** (2000), 161–230.
- [17] J.B. Zeldovich and D.A. Frank-Kamenetskii, *A theory of thermal propagation of flame*, Acta Physiochimica USSR **9** (1938), 341–350.
- [18] A. Zlatoš, *Quenching and propagation of combustion without ignition temperature cutoff*, Nonlinearity **18** (2005), 1463–1475.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN, MADISON, WI 53706, USA, EMAIL: zlatos@math.wisc.edu